

Quadratic first integrals of geodesic equations of the special type 6-dimensional h -spaces

Zolfira Zakirova[†]

Abstract

In this note we find a 6-dimensional h -spaces of the $[(21 \dots 1)(21 \dots 1) \dots (1 \dots 1)]$ type and then determine quadratic first integrals of the geodesic equations of these h -spaces.

1 Introduction

The aim of this paper is to investigate the 6-dimensional pseudo-Riemannian space $V^6(g_{ij})$ with signature $[++--]$. In particular, we find a metrics of 6-dimensional h -spaces of the $[(21 \dots 1)(21 \dots 1) \dots (1 \dots 1)]$ type and then determine quadratic first integrals of the geodesic equations of these h -spaces. The metrics of h -spaces of the $[2211]$ type has been obtained by the author [Z1]. The general method of determining pseudo-Riemannian manifolds that admit some nonhomothetic projective group G_r has been developed by A.V.Aminova [A] and has got a detailed account in the paper by the author [Z2] for the h -spaces of the type [51].

In order to find the h -spaces admitting a nonhomothetic infinitesimal projective transformation, one needs to integrate the Eisenhart equation

$$h_{ij,k} = 2g_{ij}\varphi_{,k} + g_{ik}\varphi_{,j} + g_{jk}\varphi_{,i}, \quad (1)$$

which, in the skew-normal frames, is of the form [A]

$$X_r \bar{a}_{pq} + \sum_{h=1}^n e_h (\bar{a}_{hq} \gamma_{hpr} + \bar{a}_{ph} \gamma_{hqr}) = \bar{g}_{pr} X_q \varphi + \bar{g}_{qr} X_p \varphi \quad (p, q, r = 1, \dots, n), \quad (2)$$

where

$$X_r \varphi \equiv \xi_r^i \frac{\partial \varphi}{\partial x^i}, \quad \gamma_{pqr} = -\gamma_{qpr} = \xi_{i,j} \xi_q^i \xi_r^j, \quad a_{ij} = h_{ij} - 2\varphi g_{ij},$$

[†]Kazan State University, e-mail: Zolfira.Zakirova@soros.ksu.ru, Zolfira.Zakirova@ksu.ru

ξ_i^j are the components of skew-normal frames, $\bar{g}_{pr}$ and $\bar{a}_{pq}$ are the canonical forms of the tensors g_{pr} , a_{pq} , respectively. For the type $[(21 \dots 1)(21 \dots 1) \dots (1 \dots 1)]$ h -spaces these latter have the following form [P]

$$\bar{g}_{ij}dx^i dx^j = 2e_2 dx^1 dx^2 + 2e_4 dx^3 dx^4 + \sum_{\sigma=5}^6 e_\sigma dx^{\sigma^2}, \quad (3)$$

$$\begin{aligned} \bar{a}_{ij}dx^i dx^j &= 2e_2 \lambda_2 dx^1 dx^2 + e_2 dx^{2^2} + 2e_4 \lambda_4 dx^3 dx^4 + e_4 dx^{4^2} + \sum_{\sigma=5}^6 e_\sigma \lambda_\sigma dx^{\sigma^2}, \\ (e_1 = e_2, e_3 = e_4), \quad e_i &= \pm 1, \end{aligned}$$

where $\lambda_1 = \lambda_2$, $\lambda_3 = \lambda_4$, λ_5 , λ_6 are real functions, which may coincide. These functions are roots of the characteristic equation $\det(h_{ij} - \lambda g_{ij}) = 0$.

2 The metrics of the h -spaces of the $[22(11)]$ type

Substituting the canonical forms $\bar{g}_{pr}$ and $\bar{a}_{pq}$ from (3) into (2) and taking into account that for the type $[22(11)]$ h -spaces $\lambda_5 = \lambda_6$, $\tilde{1} = 2$, $\tilde{2} = 1$, $\tilde{3} = 4$, $\tilde{4} = 3$, $\tilde{5} = 5$, $\tilde{6} = 6$, one gets the following system of equations

$$X_r \lambda_2 = 0 \ (r \neq 2), \quad X_r \lambda_4 = 0 \ (r \neq 4), \quad X_r \lambda_6 = 0, \quad (4)$$

$$\begin{aligned} X_2(\lambda_2 - \varphi) &= X_4(\lambda_4 - \varphi) = 0, \quad \gamma_{121} = e_2 X_2 \phi, \quad \gamma_{343} = e_4 X_4 \phi, \\ \gamma_{142} = \gamma_{241} &= \frac{e_2 X_4 \varphi}{\lambda_2 - \lambda_4}, \quad \gamma_{242} = -\frac{e_2 X_4 \varphi}{(\lambda_2 - \lambda_4)^2}, \quad \gamma_{324} = \gamma_{423} = \frac{e_4 X_2 \varphi}{\lambda_4 - \lambda_2}, \\ \gamma_{424} &= -\frac{e_4 X_2 \varphi}{(\lambda_4 - \lambda_2)^2}, \quad \gamma_{244} = \frac{e_4 X_2 \varphi}{(\lambda_2 - \lambda_4)^2}, \quad \gamma_{s\sigma\sigma} = \frac{e_\sigma X_s \varphi}{(\lambda_s - \lambda_\sigma)}, \end{aligned}$$

where $\sigma = 5, 6$, $s = 2, 4$, γ_{56r} are arbitrary. All others γ_{pqr} are equal to zero.

The system of linear partial differential equations

$$X_q \theta = \xi_q^i \partial_i \theta = 0, \quad (q = 1, \dots, m, i = 1, \dots, 6),$$

where ξ_q^i are the components of skew-normal frames, is completely integrable if and only if the following conditions are fulfilled (see [E] and [A])

$$(X_q, X_r) \theta = X_q X_r \theta - X_r X_q \theta = \sum_{p=1}^6 e_p (\gamma_{pqr} - \gamma_{prq}) X_{\tilde{p}}. \quad (5)$$

Using formulas (4) and (5), one can write down the commutators of the operators X_i ($i = 1, \dots, 6$) in the h -spaces under consideration,

$$(X_1, X_2) = -e_1 \gamma_{121} X_2, \quad (X_1, X_3) = 0, \quad (X_2, X_3) = e_4 \gamma_{423} X_3,$$

$$(X_1, X_4) = -e_2\gamma_{241}X_1, \quad (X_3, X_4) = -e_3\gamma_{343}X_4, \quad (6)$$

$$(X_2, X_4) = -e_2\gamma_{242}X_1 - e_1\gamma_{142}X_2 + e_4\gamma_{424}X_3 + e_3\gamma_{324}X_4,$$

$$(X_p, X_\sigma) = -e_\tau\gamma_{\tau\sigma p}X_\tau, \quad (X_q, X_\sigma) = e_\sigma\gamma_{\sigma q\sigma}X_\sigma - e_\tau\gamma_{\tau\sigma q}X_\tau,$$

$$(X_5, X_6) = -e_5\gamma_{565}X_5 + e_6\gamma_{656}X_6,$$

where $p = 1, 3, q = 2, 4, \sigma, \tau = 5, 6$ ($\sigma \neq \tau$).

After coordinate transformation $x^{i'} = \theta^i(x)$, where θ^i are solutions of the completely integrable systems from (6), one gets

$$\xi_p^i = P_p(x)\delta_p^i, \quad \xi_2^3 = \xi_2^4 = \xi_2^\sigma = \xi_4^1 = \xi_4^2 = \xi_4^\sigma = \xi_\sigma^\alpha = 0, \quad (7)$$

where $\alpha = 1, 2, 3, 4$.

Now integrating the system of equations (6) with using (4) and (7), one can find the components ξ_i^j of skew-normal frames. Using these results and the formulas [A]

$$g^{ij} = \sum_{h=1}^n e_h \xi_h^i \xi_h^j, \quad \xi_i = g_{ij} \xi_h^j, \quad a_{ij} = \sum_{h,l=1}^n e_h e_l \bar{a}_{hl} \xi_i^h \xi_j^l \quad (8)$$

we can calculate the components of tensors g_{ij} and a_{ij} . Then we come to the following

Theorem 1. *If a symmetric tensor h_{ij} of the characteristics [22(11)] and a function φ satisfy in $V^6(g_{ij})$ equation (1), then there exists a holonomic coordinate system so that the function φ and the tensors g_{ij} , h_{ij} are defined by the formulas*

$$g_{ij}dx^i dx^j = e_2 A(f_4 - f_2) \{ (f_4 - f_2) dx^1 dx^2 - A(dx^2)^2 \} + \quad (9)$$

$$e_4 \tilde{A}(f_2 - f_4) \{ (f_2 - f_4) dx^3 dx^4 - \tilde{A}(dx^4)^2 \} + F_{\sigma\tau} (f_2 - \lambda)^2 (f_4 - \lambda)^2 dx^\sigma dx^\tau,$$

$$a_{ij} dx^i dx^j = f_2 g_{i_1 j_1} dx^{i_1} dx^{j_1} + A g_{12} (dx^2)^2 + f_4 g_{i_2 j_2} dx^{i_2} dx^{j_2} + \tilde{A} g_{34} (dx^4)^2 + \lambda g_{\sigma\tau} dx^\sigma dx^\tau, \quad (10)$$

$$h_{ij} = a_{ij} + 2\varphi g_{ij}, \quad 2\varphi = 2f_2 + 2f_4 + c, \quad (11)$$

$$A = \epsilon x^1 + \theta(x^2), \quad \tilde{A} = \tilde{\epsilon} x^3 + \omega(x^4), \quad (12)$$

where $\epsilon, \tilde{\epsilon} = 0, 1$, $f_2 = \epsilon x^2$, $f_4 = \tilde{\epsilon} x^4 + a$, λ, c and a are some constants, $a \neq 0$ when $\tilde{\epsilon} = 0$, $F_{\sigma\tau}(x^5, x^6), \theta(x^2), \omega(x^4)$ are arbitrary functions, $\theta \neq 0$ when $\epsilon = 0$, $\omega \neq 0$ when $\tilde{\epsilon} = 0$, $i_1, j_1 = 1, 2$, $i_2, j_2 = 3, 4$, $\sigma, \tau = 5, 6$, $e_2, e_4 = \pm 1$.

3 The metrics of the h -spaces of the $[2(21)1]$, $[2(211)]$ types

Substituting the canonical forms $\bar{g}_{pr}$ and $\bar{a}_{pq}$ from (3) into (2) and taking into account that for the type $[2(21)1]$ h -spaces $\lambda_4 = \lambda_5$, one gets the following system of equations

$$X_r \lambda_2 = 0 \ (r \neq 2), \quad X_r \lambda_5 = 0, \quad X_r \lambda_6 = 0 \ (r \neq 6), \quad (13)$$

$$\begin{aligned} X_2(\lambda_2 - \varphi) = X_6(\lambda_6 - \varphi) = 0, \quad \gamma_{121} = e_2 X_2 \phi, \\ \gamma_{162} = \gamma_{261} = \frac{e_2 X_6 \varphi}{\lambda_2 - \lambda_6}, \quad \gamma_{262} = -\frac{e_2 X_6 \varphi}{(\lambda_2 - \lambda_6)^2}, \quad \gamma_{3s4} = \gamma_{4s3} = \frac{e_4 X_s \varphi}{\lambda_5 - \lambda_s}, \\ \gamma_{4s4} = -\frac{e_4 X_s \varphi}{(\lambda_5 - \lambda_s)^2}, \quad \gamma_{2\sigma\sigma} = \frac{e_\sigma X_2 \varphi}{(\lambda_2 - \lambda_\sigma)}, \quad \gamma_{565} = \frac{e_5 X_6 \varphi}{\lambda_5 - \lambda_6}, \end{aligned}$$

where $\sigma = 5, 6$, $s = 2, 6$, γ_{45r} are arbitrary, all others γ_{pqr} are equal to zero.

The commutators of the operators X_i ($i = 1, \dots, 6$) in the h -spaces of the $[2(21)1]$ type have the form

$$(X_1, X_2) = -e_1 \gamma_{121} X_2, \quad (X_1, X_3) = 0, \quad (X_2, X_3) = e_4 \gamma_{423} X_3, \quad (14)$$

$$(X_1, X_4) = -e_5 \gamma_{541} X_5, \quad (X_1, X_5) = -e_4 \gamma_{451} X_3, \quad (X_1, X_6) = -e_2 \gamma_{261} X_1,$$

$$(X_2, X_4) = e_3 \gamma_{324} X_4 + e_4 \gamma_{424} X_3 - e_5 \gamma_{542} X_5, \quad (X_2, X_5) = e_5 \gamma_{525} X_5 - e_4 \gamma_{425} X_3,$$

$$(X_2, X_6) = -e_2 \gamma_{262} X_1 - e_1 \gamma_{162} X_2 + e_6 \gamma_{626} X_6, \quad (X_3, X_4) = -e_5 \gamma_{543} X_5,$$

$$(X_3, X_5) = -e_4 \gamma_{453} X_3, \quad (X_3, X_6) = -e_4 \gamma_{463} X_3, \quad (X_4, X_5) = -e_4 \gamma_{454} X_3 + e_5 \gamma_{545} X_5,$$

$$(X_4, X_6) = -e_3 \gamma_{364} X_4 - e_4 \gamma_{464} X_3 + e_5 \gamma_{546} X_5, \quad (X_5, X_6) = e_4 \gamma_{456} X_3 - e_5 \gamma_{565} X_5.$$

The systems $X_i \theta = 0$ ($i \neq 2$), $X_j \theta = 0$ ($j \neq 4$), $X_k \theta = 0$ ($k \neq 6$) are completely integrable and have the following solutions: θ^2 , θ^4 , θ^6 . The systems $X_3 \theta = X_4 \theta = X_5 \theta = X_6 \theta = 0$, $X_1 \theta = X_2 \theta = X_3 \theta = X_6 \theta = 0$ and $X_1 \theta = X_2 \theta = X_6 \theta = 0$ are completely integrable too. The first system has the solutions θ^1 and θ^2 , the second system has the solutions θ^4 and θ^5 , the third system has the solutions θ^3 , θ^4 and θ^5 . After coordinate transformation $x^{i'} = \theta^i(x)$, one gets

$$\xi_p^i = P_p(x) \delta_p^i, \quad \xi_2^s = \xi_4^q = \xi_5^q = \xi_5^4 = 0, \quad (15)$$

where $p = 1, 2, 3$, $s = 3, 4, 5, 6$, $q = 1, 2, 6$.

Now integrating the system of equations (14) with using (13) and (15), and then calculating the components of tensors g_{ij} and a_{ij} we come to the following results

$$g_{ij}dx^i dx^j = e_2\{2(f_6 - f_2)Adx^1 dx^2 - A^2(dx^2)^2\} + \quad (16)$$

$$(f_6 - \lambda)(f_2 - \lambda)^2\{2e_4 dx^3 dx^4 - e_4(\Sigma + \omega)(dx^4)^2 + e_5(dx^5)^2\} + e_6(f_2 - f_6)^2(dx^6)^2, \\ a_{ij}dx^i dx^j = f_2 g_{i_1 j_1} dx^{i_1} dx^{j_1} + g_{12}(dx^2)^2 + \lambda g_{i_2 j_2} dx^{i_2} dx^{j_2} + g_{34}(dx^4)^2 + f_6 g_{66}(dx^6)^2, \quad (17)$$

$$h_{ij} = a_{ij} + (2f_2 + f_6 + c)g_{ij}, \quad \varphi = f_2 + \frac{1}{2}f_6 + c, \quad (18)$$

$$A = \epsilon x^1 + \theta(x^2), \quad \Sigma = 2(f_2 - \lambda)^{-1} + (f_6 - \lambda)^{-1}, \quad (19)$$

where $\epsilon = 0, 1$, $f_2 = \epsilon x^2$, λ and c are some constants, $\theta(x^2), \omega(x^4, x^5), f_6(x^6)$ are arbitrary function, $\theta \neq 0$ when $\epsilon = 0$, $i_1, j_1 = 1, 2$, $i_2, j_2 = 3, 4, 5$, $e_2, e_4, e_5, e_6 = \pm 1$.

After same computations for h -spaces of the $[2(211)]$ type, we obtain

$$g_{ij}dx^i dx^j = 2e_2 Adx^1 dx^2 + (f_2 - \lambda)^2\{2e_4 dx^3 dx^4 - e_4(\Sigma + \omega)(dx^4)^2 + g_{\sigma\tau} dx^\sigma dx^\tau\}, \quad (20)$$

$$a_{ij}dx^i dx^j = 2f_2 g_{12} dx^1 dx^2 + g_{12}(dx^2)^2 + \lambda g_{pq} dx^p dx^q + g_{34}(dx^4)^2, \quad (21)$$

$$h_{ij} = a_{ij} + (2f_2 + c)g_{ij}, \quad \varphi = f_2 + c, \quad (22)$$

$$A = \epsilon x^1 + \theta(x^2), \quad \Sigma = 2(f_2 - \lambda)^{-1}, \quad (23)$$

where $\epsilon = 0, 1$, $f_2 = \epsilon x^2$, λ, c are some constants, $\theta(x^2), \omega(x^4, x^5, x^6), g_{\sigma\tau}(x^4, x^5, x^6)$ are arbitrary function, $\theta \neq 0$ when $\epsilon = 0$, $p, q = 3, 4, 5, 6$, $\sigma, \tau = 5, 6$, $e_2, e_4 = \pm 1$.

Let us formulate these results in following theorem.

Theorem 2. *If a symmetric tensor h_{ij} of the characteristics $[2(21)1]$, $[2(211)]$ and a function φ satisfy in $V^6(g_{ij})$ the Eisenhart equation, then there exists a holonomic coordinate system so that the function φ and the tensors g_{ij} , h_{ij} are defined by formulas (16)-(23).*

4 The metrics of the h -spaces of the $[(22)11]$,

$[(221)1]$ types

For the h -spaces of the $[(22)11]$ type from (2), we get the following system of equations

$$X_r \lambda_4 = 0, \quad X_r \lambda_\sigma = 0 \ (r \neq \sigma), \quad X_\sigma(\lambda_\sigma - \varphi) = 0,$$

$$\begin{aligned}\gamma_{14r} = \gamma_{23r}, \quad \gamma_{1\sigma 2} = \gamma_{2\sigma 1} = \frac{e_2 X_\sigma \varphi}{\lambda_4 - \lambda_\sigma}, \quad \gamma_{3\sigma 4} = \gamma_{4\sigma 3} = \frac{e_4 X_\sigma \varphi}{\lambda_4 - \lambda_\sigma}, \\ \gamma_{s\sigma s} = -\frac{e_s X_\sigma \varphi}{(\lambda_4 - \lambda_\sigma)^2}, \quad \gamma_{\sigma\tau\sigma} = \frac{e_\sigma X_\tau \varphi}{\lambda_\sigma - \lambda_\tau} \quad (\sigma \neq \tau),\end{aligned}\tag{24}$$

where $\sigma, \tau = 5, 6$, $s = 2, 4$, γ_{24r} are arbitrary, while all others γ_{pqr} are equal to zero.

In this case the system of equations in components ξ_i^j of skew-normal frames has the form

$$\begin{aligned}1^\circ \quad \xi_1^\alpha \partial_\alpha \xi_2^\beta - \xi_2^\alpha \partial_\alpha \xi_1^\beta &= \gamma_{412} \xi_3^\beta - \gamma_{421} \xi_3^\beta - \gamma_{411} \xi_4^\beta, \\ 2^\circ \quad \xi_1^\alpha \partial_\alpha \xi_3^\beta - \xi_3^\alpha \partial_\alpha \xi_1^\beta &= \gamma_{413} \xi_3^\beta - \gamma_{141} \xi_1^\beta, \\ 3^\circ \quad \xi_1^\alpha \partial_\alpha \xi_4^\beta - \xi_4^\alpha \partial_\alpha \xi_1^\beta &= -\gamma_{141} \xi_2^\beta - \gamma_{414} \xi_3^\beta - \gamma_{241} \xi_1^\beta, \\ 4^\circ \quad \xi_\sigma^\sigma \partial_\sigma \xi_1^\beta &= \frac{1}{2} \frac{f'_\sigma}{\lambda - f_\sigma} \xi_\sigma^\sigma \xi_1^\beta - \gamma_{41\sigma} \xi_3^\beta, \\ 5^\circ \quad \xi_2^\alpha \partial_\alpha \xi_3^p - \xi_3^\alpha \partial_\alpha \xi_2^p &= -\gamma_{142} \xi_1^p + \gamma_{413} \xi_4^p + \gamma_{423} \xi_3^p, \\ 6^\circ \quad \xi_3^\alpha \partial_\alpha \xi_2^q &= -\gamma_{413} \xi_4^q, \\ 7^\circ \quad \xi_2^\alpha \partial_\alpha \xi_4^\beta - \xi_4^\alpha \partial_\alpha \xi_2^\beta &= -\gamma_{142} \xi_2^\beta + \gamma_{414} \xi_4^\beta - \gamma_{242} \xi_1^\beta + \gamma_{424} \xi_3^\beta, \\ 8^\circ \quad \xi_\sigma^\sigma \partial_\sigma \xi_2^\beta &= \frac{1}{2} \frac{f'_\sigma}{\lambda - f_\sigma} \xi_\sigma^\sigma \xi_2^\beta - \frac{1}{2} \frac{f'_\sigma}{(\lambda - f_\sigma)^2} \xi_\sigma^\sigma \xi_1^\beta - \gamma_{41\sigma} \xi_4^\beta - \gamma_{42\sigma} \xi_3^\beta, \\ 9^\circ \quad \xi_3^\alpha \partial_\alpha \xi_4^\beta - \xi_4^\alpha \partial_\alpha \xi_3^\beta &= -\gamma_{143} \xi_2^\beta + \gamma_{144} \xi_1^\beta - \gamma_{243} \xi_1^\beta, \\ 10^\circ \quad \xi_\sigma^\sigma \partial_\sigma \xi_3^\beta &= \frac{1}{2} \frac{f'_\sigma}{\lambda - f_\sigma} \xi_\sigma^\sigma \xi_3^\beta - \gamma_{14\sigma} \xi_1^\beta, \\ 11^\circ \quad \xi_\sigma^\sigma \partial_\sigma \xi_4^\beta &= \frac{1}{2} \frac{f'_\sigma}{\lambda - f_\sigma} \xi_\sigma^\sigma \xi_4^\beta - \frac{1}{2} \frac{f'_\sigma}{(\lambda - f_\sigma)^2} \xi_\sigma^\sigma \xi_3^\beta - \gamma_{14\sigma} \xi_2^\beta - \gamma_{24\sigma} \xi_1^\beta, \\ 12^\circ \quad \xi_\beta^\alpha \partial_\alpha \xi_\sigma^\sigma &= 0, \\ 13^\circ \quad \xi_\sigma^\sigma \partial_\sigma \xi_\tau^\tau &= \frac{1}{2} \frac{f'_\sigma}{f_\tau - f_\sigma} \xi_\sigma^\sigma \xi_\tau^\tau, \quad (\tau \neq \sigma),\end{aligned}$$

where $\alpha, \beta = 1, 2, 3, 4$, $p = 1, 3$, $q = 2, 4$, $\sigma, \tau = 5, 6$, $\xi_\sigma^i = P_\sigma(x) \delta_\sigma^i$, $\xi_p^q = \xi_\alpha^\sigma = 0$.

Integrating 12° and 13°, after coordinate transformation x^5, x^6 one gets

$$\xi_5^5 = (f_6 - f_5)^{-1/2}, \quad \xi_6^6 = (f_5 - f_6)^{-1/2}.$$

Differentiating $g^{\alpha\beta} = \sum_{h=1}^6 e_h \xi_h^\alpha \xi_h^\beta$ with respect to x^p , with using 1° – 3° and 5° – 7°, one gets $\partial_p g^{\alpha\beta} = 0$. Differentiating $g^{\alpha\beta}$ with respect to x^σ , from equations 4°, 8°, 10°, 11°, one obtains

$$\partial_\sigma g^{pr} = -\frac{f'_\sigma}{f_\sigma - \lambda} g^{pr} - \frac{f'_\sigma}{(f_\sigma - \lambda)^2} \Pi_\sigma (f_\sigma - \lambda)^{-1}, \quad \partial_\sigma g^{pq} = -\frac{f'_\sigma}{f_\sigma - \lambda} g^{pq},$$

where $p, r = 1, 3$. Integrating this equations, one gets

$$g^{pr} = \Pi_\sigma (f_\sigma - \lambda)^{-1} (\Sigma_1 + F^{pr}), \quad g^{pq} = \Pi_\sigma (f_\sigma - \lambda)^{-1} F^{pq},$$

where F^{pr}, F^{pq} are functions of the variables x^2, x^4 .

After same calculations for the h -spaces of the $[(221)1]$ type one obtains

$$g_{ij}dx^i dx^j = \Pi_\sigma(f_\sigma - \lambda)\{2g_{12}dx^1 dx^2 - e_2(\Sigma_1 + \theta_1)(dx^2)^2 + \quad (25)$$

$$2g_{34}dx^3 dx^4 - e_4(\Sigma_1 + \theta_2)(dx^4)^2 + G\} + \sum_\sigma e_\sigma(f_\tau - f_\sigma)(dx^\sigma)^2, \\ a_{ij}dx^i dx^j = \lambda(g_{st}dx^s dx^t + G) + g_{12}(dx^2)^2 + g_{34}(dx^4)^2 + \sum_\sigma f_\sigma g_{\sigma\sigma}(dx^\sigma)^2 + G, \quad (26)$$

$$h_{ij} = a_{ij} + \left(\sum_\sigma f_\sigma + c\right)g_{ij}, \quad \varphi = \frac{kf_k}{2} + \frac{1}{2}\sum_\sigma f_\sigma + c. \quad (27)$$

$$\Sigma_1 = \sum_\sigma (f_\sigma - \lambda)^{-1}, \quad (28)$$

$$G = 2e_5\{1 + \theta_3(f - \lambda)\}dx^4 dx^5 + (f - \lambda)g_{55}(dx^5)^2 + e_6(dx^6)^2, \quad (29)$$

where λ, c are some constants. Here for the h -spaces of the $[(22)11]$ type $\tau, \sigma = 5, 6$ ($\tau \neq \sigma$), $s, t = 1, 2, 3, 4$, $G = 0$, $g_{st}, \theta_1, \theta_2$ are functions of the variables x^2, x^4 , f_σ are functions of the variable x^σ . For the h -spaces of the $[(221)1]$ type $\sigma = 6$, $s, t = 1, 2, 3, 4, 5$, $g_{st}, \theta_1, \theta_2, \theta_3$ are functions of the variables x^2, x^4, x^5 , f_6 is a function of the variable x^6 .

Theorem 3. *If a symmetric tensor h_{ij} of the characteristics $[(22)11]$, $[(221)1]$ and a function φ satisfy in $V^6(g_{ij})$ the Eisenhart equation, then there exists a holonomic coordinate system so that the function φ and the tensors g_{ij} , h_{ij} are defined by formulas (25)-(29).*

5 The metrics of the h -spaces of the $[(2211)]$, $[(22)(11)]$ and $[(21)(21)]$ types

In this cases, the function φ is constant, therefore, from (1) the tensor h_{ij} is covariant constant. We skip further calculations that are just integration of the equations with respect to ξ_i^j combined with some proper chosen coordinate transformations. The result reads

for the h -spaces of the $[(2211)]$ type

$$g_{ij}dx^i dx^j = 2g_{12}dx^1 dx^2 - e_2\theta(dx^2)^2 + 2g_{34}dx^3 dx^4 + g_{rq}dx^r dx^q, \quad (30)$$

$$a_{ij}dx^i dx^j = \lambda g_{ij}dx^i dx^j + g_{12}(dx^2)^2, \quad (31)$$

$$h_{ij} = a_{ij} + c g_{ij}, \quad (32)$$

where $r, q = 5, 6$, λ, c are some constants, $\theta, g_{12}, g_{34}, g_{rq}$ are functions of the variables x^2, x^4, x^5, x^6 ,

for the h -spaces of the $[(22)(11)]$ type

$$g_{ij}dx^i dx^j = e_2\{2dx^1 dx^2 - \theta(dx^2)^2\} + e_4\{2dx^3 dx^4 - \omega(dx^4)^2\} + g_{\sigma\tau}dx^\sigma dx^\tau, \quad (33)$$

$$a_{ij}dx^i dx^j = \lambda_1\{g_{i_1 j_1} dx^{i_1} dx^{j_1} + e_2(dx^2)^2 + g_{i_2 j_2} dx^{i_2} dx^{j_2} + e_4(dx^4)^2\} + \lambda_2 g_{\sigma\tau} dx^\sigma dx^\tau, \quad (34)$$

$$h_{ij} = a_{ij} + c g_{ij}, \quad (35)$$

where θ, ω are functions of the variables x^2, x^4 , $g_{\sigma\tau}$ are functions of the variables x^5, x^6 , λ_1, λ_2, c are some constants, here $\lambda_1 \neq \lambda_2$, $i_1, j_1 = 1, 2$, $i_2, j_2 = 3, 4$, $\sigma, \tau = 5, 6$,

for the h -spaces of the $[(21)(21)]$ type

$$g_{ij}dx^i dx^j = e_2\{2dx^1 dx^2 - \theta(dx^2)^2\} + e_3(dx^3)^2 + e_5\{2dx^4 dx^5 - \omega(dx^5)^2\} + e_6(dx^6)^2, \quad (36)$$

$$a_{ij}dx^i dx^j = \lambda_1 g_{i_1 j_1} dx^{i_1} dx^{j_1} + e_2(dx^2)^2 + \lambda_2 g_{i_2 j_2} dx^{i_2} dx^{j_2} + e_5(dx^5)^2, \quad (37)$$

$$h_{ij} = a_{ij} + c g_{ij}, \quad (38)$$

where θ is a function of the variables x^2, x^3 , ω is a function of the variables x^5, x^6 , λ_1, λ_2, c are some constants, $\lambda_1 \neq \lambda_2$, $i_1, j_1 = 1, 2, 3$, $i_2, j_2 = 4, 5, 6$.

Now we have the following theorem.

Theorem 4. *If a symmetric tensor h_{ij} of the characteristics $[(2211)]$, $[(22)(11)]$, $[(21)(21)]$ and a function φ satisfy in $V^6(g_{ij})$ equation (1), then there exists a holonomic coordinate system so that the function φ and the tensors g_{ij} , h_{ij} are defined by formulas (30)-(38).*

6 Quadratic first integrals of the geodesic

equation of the h -spaces of the $[(21 \dots 1)(21 \dots 1) \dots (1 \dots 1)]$ type

For every solution h_{ij} of equation (1) there is a quadratic first integral of the geodesic equations (see [A], [Z2])

$$(h_{ij} - 4\varphi g_{ij})\dot{x}^i \dot{x}^j = \text{const}, \quad (39)$$

where $\dot{x}^i$ is the tangent vector to the geodesic.

Therefore, the quadratic first integrals of the geodesic equations of the h -spaces of the $[(21 \dots 1)(21 \dots 1) \dots (1 \dots 1)]$ type are determined by formula (39), where the tensors h_{ij} , g_{ij} and the function φ are obtained in Theorems 1-4.

I am grateful to professor A.V.Aminova for constant attention to this work and for useful discussions. The research was partially supported by the RFBR grant 01-02-17682-a and by the INTAS grant 00-00334.

References

- [A] Aminova A. V. Lie algebras of infinitesimal projective transformations of Lorentz manifolds (in Russian)// Uspekhi Mat. Nauk **50** (1995), no. 1(301), 69-142.
- [E] Eyzenhart L. P. Riemannian geometry// M.:IL, 1948.
- [P] Petrov A. Z. On geodesic mapping of the Riemannian space with indefinite metric (in Russian)// Uch. zap. Kazan. un-ta. 1949. T. 109 no. 3. 7-36.
- [Z1] Zakirova Z. Kh. 6-dimensional h -spaces of the special type (in Russian)// Lobachevskiy Int. geom. seminar, Kazan, 1997: Abstracts of contributed papers – Kazan, 1997.- P. 52.
- [Z2] Zakirova Z. Kh. First integral of the geodesics equations in 6D h -spaces (in Russian)// Proc. of Geometry Seminar, Kazan, 1997, N23. P. 57-64.